

51

BUNDESREPUBLIK DEUTSCHLAND

DEUTSCHES PATENTAMT



52

Int. Cl.:

C 07 c

C 07 d

A 01 n, 9/20

Deutsche Kl.:

12 <sup>9</sup>/<sub>21</sub>

12 o, 21

12 o, 25

12 qu, 24

12 p, 10/01

451, 9/20

10

11

21

22

43

# Offenlegungsschrift 1914 954

Aktenzeichen: P 19 14 954.3

Anmeldetag: 24. März 1969

Offenlegungstag: 23. Oktober 1969

Ausstellungspriorität: —

30

Unionspriorität

32

Datum:

25. März 1968

14. Februar 1969

33

Land:

Großbritannien

31

Aktenzeichen:

14198-68

14198

54

Bezeichnung:

Fungizide Mittel mit einem Gehalt an Acrylamidverbindungen, neue Acrylamidverbindungen und Verfahren zu ihrer Herstellung

61

Zusatz zu:

—

62

Ausscheidung aus:

—

71

Anmelder:

Shell Internationale Research Maatschappij N. V., Den Haag

Vertreter:

Wuesthoff, Dr.-Ing. F.; Puls, Dipl.-Ing. G.; von Pechmann, Dr. E.; Behrens, Dr.-Ing. D.; Patentanwälte, 8000 München

72

Als Erfinder benannt:

Haken, Pieter Ten, Herne Bay, Kent (Großbritannien)

Benachrichtigung gemäß Art. 7 § 1 Abs. 2 Nr. 1 d. Ges. v. 4. 9. 1967 (BGBl. I S. 960): —

vgl. Ber. - L. 47/73

DT 1914954

L4 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2007 ACS on STN  
 ACCESSION NUMBER: 1970:31613 CAPLUS <<LOGINID::20070925>>  
 DOCUMENT NUMBER: 72:31613  
 TITLE: Acrylamide-based fungicides  
 INVENTOR(S): Haken, Pieter T.  
 PATENT ASSIGNEE(S): Shell Internationale Research Maatschappij N. V.  
 SOURCE: Ger. Offen., 38 pp.  
 CODEN: GWXXBX  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 1914954	A	19691023	DE 1969-1914954	19690324
DE 1914954	B2	19770818		
DE 1914954	C3	19791025		
GB 1215066	A	19701209	GB 1968-14198	19680325
NL 169183	B	19820118	NL 1968-14198	19680325
BE 730347	A	19690924	BE 1969-730347	19690324
NL 6904465	A	19690929	NL 1969-4465	19690324
NL 169183	B	19820118		
NL 169183	C	19870116		
FR 2004656	A5	19691128	FR 1969-8564	19690324
CH 507650	A	19710531	CH 1969-507650	19690324
DK 125505	B	19730305	DK 1969-1603	19690324
IL 31889	A	19740910	IL 1969-31889	19690324
SE 385262	B	19760621	SE 1969-4059	19690324
SE 416808	B	19810209	SE 1973-11762	19730829
SE 416808	C	19810521		
PRIORITY APPLN. INFO.:			GB 1968-14198	A 19680325
			GB 1969-14198	A 19690214

GI For diagram(s), see printed CA Issue.

AB Fungus diseases of growing plants can be controlled by the title compds. A mixture of 4 g 2-methyl-4,5-dihydrofuran-3-carboxylic acid and 20 ml SO cl2 was heated 2 hr to give residue which was dissolved in 50 ml dry benzene and mixed with a solution of 6.5 g PhNH2 in 25 ml benzene. The mixture was kept 2 days to give I, m. 82-4° (petroleum ether). A mixture of 2-methyl-5,6-dihydro-(4H)-pyran-3-carboxylic acid, PhN:PNHPh, and PhMe was heated 4 hr to yield II, m. 109-10°. 2-Methyl-5,6-dihydro-(4H)-thiapyran-3-carboxanilide, m. 150°, prepared as above, was oxidized with H2O2 in HOAc to give III, m. 225°. A Grignard reaction using 7.2 g Mg, 42.6 g MeI, 32.1 g PhNHMe and 18.9 g 2-methyl-3-ethoxycarbonyl-5,6-dihydro-(4H)-pyran (IIIa) in ether was employed to prepare IV, m. 63-5°. NaH (1.8 g), 8.1 g PhNHNH2 and 100 ml dry tetrahydrofuran were stirred 2 hr at 20°, 8.5 g IIIa was added and the mixture was stirred 0.5 hr at 20° and 1.25 hr at reflux temperature to give V, m. 108-10° (benzene-cyclohexane). The following compds. were similarly prepared: 2-methylfuran-3-carboxanilide, m. 109-11°; 2-methylfuran-3-(N-3-tolylcarboxamide), m. 89-91°; 4-methyl-1,3-oxazole-5-carboxanilide, m. 117-18°; 2-methylcyclohex-1-ene-1-carboxanilide, m. 112-16°; 2-methylcyclopent-1-ene-1-carboxanilide, m. 93-4°; 3-(2-furyl)-crotonanilide, m. 140-1°; 3-propylcrotonanilide, m. 63.5-4.0°; 3-trifluoromethylcrotonanilide, m. 100-3°; 3-cyclopropylcroton-anilide, m. 84.5-6.5°; 3-(3-thienyl)crotonanilide, m. 160-2°; 5-methyl-1,2,3-thiadiazole-4-carboxanilide, m. 147°; 5-methyl-1,2,3-triazole-4-carboxanilide, m. 197°; 2-methyl-5,6-dihydro-(4H)-pyran-3-(N-2-biphenylcarboxamide), m. 79-81°; 3-methylpyridine-2-carboxanilide, m. 72°;

2-methyl-5,6-dihydro-(4H)-pyran-3-(N-methoxyphenyl)carboxamide ( ), m. 157°; 2-methyl-5,6-dihydro-(4H)-pyran-3-(N-cyclohexylcarboxamide), m. 154°; 2-methylbenzo[b]furan-3-carboxanilide, m. 143-5°; 2-ethyl-5,6-dihydro-(4H)-pyran-3-carboxanilide, m. 119°; 2-methyl-5,6-dihydro-(4H)-pyran-3-[N-(4-fluorophenyl)carboxamide], m. 133-5°; 2-methyl-5,6-dihydro-(4H)-pyran-3-[N-(4-ethoxyphenyl)carboxamide], m. 145-8°; 2,3-dimethylcrotonanilide, m. 92-3°; 2-methyl-5,6-dihydro-(4H)-pyran-3-[N-(4,5-methylenedioxyphenyl)carboxamide], m. 127-9°; 2,6-dimethyl-5,6-dihydro-(4H)-pyran-3-carboxanilide, m. 112-13°; 2-methylfuran-3-[N-(4-methoxyphenyl)carboxamide], m. 104-5°; 2-methyl-5,6-dihydro-(4H)-pyran-3-[(N-(2-methoxyphenyl)carboxamide], m. 98-9°; 2-methylfuran-3-(N-cyclohexylcarboxamide), m. 99-100°; 2-methyl-5,6-dihydro-(4H)-pyran-3-[N-(3-methoxyphenyl)carboxamide], m. 104-6°; 3-cyclohex-1-enylcrotonanilide, m. 152-3°. Bean rust and cucumber mildew were controlled by applications of selected compds. to seeds, roots or foliage in concns. of 100-5000 ppm.

IT 24691-92-7P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 24691-92-7 CAPLUS

CN v-Triazole-4-carboxanilide, 5-methyl- (8CI) (CA INDEX NAME)

